

A Comparative Study on Short-term Clinical Outcome in Acute Coronary Syndrome

Sultana SA¹, Iqbal SA², Malek MA³, Rahman MW⁴, Alam AYMS⁵, Haque MA⁶, Rabbani SB⁷

Abstract

Introduction: Coronary artery diseases are currently the major cause of death in developing countries. Acute coronary syndrome (ACS) is defined as any group of clinical symptoms compatible with acute myocardial ischemia and covers the spectrum of clinical conditions ranging from unstable angina (UA) to non-ST elevation myocardial infarction (NSTEMI) to ST-elevation myocardial infarction (STEMI). Accurate diagnosis and management of ACS has life-saving implications of its outcome.

Objective: To compare the outcomes of STEMI and NSTEMI in a percutaneous coronary intervention (PCI) capable centre.

Materials and Methods: The patients who undergone percutaneous coronary intervention (PCI) in Combined Military Hospital, Dhaka were considered from January 2013 to January 2017. Diagnosis of acute MI was based on the clinical presentation, electrocardiogram (ECG) and raised highly sensitive troponin I. Acute MI patients were classified into 2 groups, STEMI and NSTEMI. Their coronary risk factors, co-morbidity, ECG, echocardiogram, coronary angiographic (CAG) findings and short-term outcomes were collected. All statistical data were analysed by SPSS 22.0 software.

Results: There were 464 patients enrolled for analysis. Among them, 208(44.8%) patients had STEMI and 256(55.2%) had NSTEMI. The ratio of male/female was greater in STEMI as compared to NSTEMI (4.0 vs 1.9; $p=0.041$). Among NSTEMI patients, 88(34.4%) had ST depression, 168(65.6%) patients had other ECG changes like T wave abnormalities in 66(25.7%) and poor R-wave progression in 16(6.3%). NSTEMI patients had less regional wall motion abnormality on echo cardiogram ($p=0.0045$). As a complication heart failure (36% vs 9.3%), cardiogenic shock (16.8% vs 15.6%), atrial fibrillation (7.2 vs 0.78 %), ventricular tachycardia (2.8% vs 0.5%), reinfarction (3 % vs 0.78%) and death (2.4% vs 0.40%) were observed more in STEMI patients than NSTEMI respectively. NSTEMI patients had

less regional wall motion abnormality on echocardiogram ($p=0.0045$). As a complication heart failure (36% vs 9.3%), cardiogenic shock (16.8% vs 15.6%), atrial fibrillation (7.2% vs 0.78%), ventricular tachycardia (2.8% vs 0.5%), reinfarction (3% vs 0.78%) and death (2.4% vs 0.40%) were observed more in STEMI patients than NSTEMI respectively. Coronary angiogram shows that left anterior descending artery was the most commonly involved artery in STEMI; however, the left circumflex artery or right coronary artery was involved more commonly in NSTEMI ($p<0.001$).

Conclusion: The first step in successful treatment of acute MI depends on early diagnosis. In spite of immediate management, STEMI had relatively worse outcome compared to NSTEMI.

Key-words: Acute coronary artery disease, Acute coronary syndrome, ST-elevation myocardial infarction (STEMI), Non-ST elevation myocardial infarction (NSTEMI).

Introduction

Acute myocardial infarction (MI) significantly contributes to mortality and morbidity in developing countries. To reduce mortality and morbidity, early clinical diagnosis of obstructive coronary artery diseases with subsequent management is necessary for all patients with chest discomfort or other symptoms suggestive of an acute coronary syndrome (ACS)¹⁻³. Although there are increasing numbers of methods to diagnose acute MI, but electrocardiography (ECG) is considered a quick, easily accessible tool, and it is the most important initial investigation for diagnosis of myocardial ischemia and MI⁴. ST-segment elevation MI (STEMI) can easily be diagnosed from clinical manifestations and typical ECG changes. The introduction of new therapeutic modalities, including invasive cardiac procedures and new medications, play a major role in outcome of these patients. We conducted this study to compare outcome of NSTEMI and STEMI patients.

1. **Col Syeda Aleya Sultana**, MBBS, D CARD (China), DAM, FCPS, FACC (USA), Classified Specialist in Cardiology, CMH, Dhaka
 2. **Brig General Syed Asif Iqbal**, MBBS, FCPS(Med), FACC (USA), FRCP, Chief Physician, CMH, Dhaka
 3. **Col Md. Abdul Malek**, MBBS, MCPS(MED), FCPS(MED), FCPS (CARD), Classified Specialist in Cardiology, CMH, Dhaka
 4. **Lt Col Md. Wali-ur-Rahman**, FCPS(Med), FCPS (Card) Classified Spl in Medicine (Cardiology), CMH, Dhaka
 5. **Lt Col Abu Yusuf Md. Shahidul Alam**, MBBS, D Card, FCPS (Med), Classified Specialist in Medicine (Cardiology), CMH, Dhaka
 6. **Maj Md. Aminul Haque**, MBBS, MD(Card) Classified Spl in Medicine (Cardiology), CMH, Dhaka
 7. **Dr Safia Binte Rabbani**, MBBS, MCPS(Med), FCPS(Med), Trainee in cardiology, CMH, Dhaka.

Materials and Methods

This retrospective study was carried out in Combined Military Hospital, Dhaka from January 2013 to January 2017 with a view to compare short term clinical outcome between STEMI and NSTEMI patients. There were 464 patients enrolled for analysis and among them 208 patients (44.8%) had STEMI and 256(55.2%) had NSTEMI. The diagnostic criteria of STEMI were an ST segment elevation of ≥ 2 mm in adjacent chest leads and/or an ST segment elevation of >1 mm in two or more standard limb leads or a new Left Bundle Branch Block (LBBB) and presence of positive cardiac biomarkers. NSTEMI was diagnosed in the absence of ST segment elevation and positive highly sensitive troponin I, CK-MB. PCI was performed according to standard procedures. Their age, coronary risk factors, co-morbidities, ECG, echocardiogram, angiographic findings and short-term outcomes were collected. All statistical data were analyzed by SPSS 22.0 software and $p < 0.05$ was considered statistically significant.

Results

Among 464 patients, 256 were included in NSTEMI and 208 patients from STEMI group; 164(78.84%) male patients sustained acute STEMI whereas female were 44(21.15%). In NSTEMI male patients were 172(67.1%) and female were 84(32.8%).

Distribution of patients according to sex, age, cardiac risk factors are shown in Figure-1,2,3 respectively. Compared to patients of STEMI, patients of NSTEMI were older. The incidence of cardiovascular risk factors such as hypertension, diabetes and hypercholesterolemia were higher in NSTEMI patients than patients of STEMI ($P < 0.05$). In addition, coronary angiographic (CAG) shows that the incidence of multivessel involvement in NSTEMI patients were higher than that in STEMI patients ($P < 0.001$).




Fig-1: Patient distribution in MI




Fig-2: Age distribution among MI patients




Fig-3: Major CAD risk factors for MI patients

Identification of ST-segment elevation in ECGs is easy; however, NSTEMI patients had various types of ECG findings (Figure-4). Among NSTEMI patients, 88 (34.4%) had ST depression that may have been true ST depression or reciprocal changes. 120 patients had other ECG changes, including T-wave abnormalities (n=66; 25.7%), poor R-wave progression (n=16; 6.3%), atrial fibrillation (n=12; 4.7%), bundle branch block (n=4; 1.5%), frequent premature ventricular contraction (n=2; 0.7%).




Fig-4: ECG changes in the corresponding lead in NSTEMI patients

Table-I: Echocardiographic and CAG findings

| | Title | STEMI, n (%) | NSTEMI, n (%) | P value |
|---------------------------------|------------------------|---------------------|----------------------|----------------|
| Wall motion on echocardiography | No RWMA | 08 (3.9) | 88 (34.4) | 0.0045 |
| | Hypokinesia | 32 (15.5) | 94 (36.7) | |
| | Akinesia | 152 (73.8) | 62 (24.2) | |
| | Dyskinesia | 14 (6.8) | 12 (04.7) | |
| LV function on echocardiography | Fair ($\geq 55\%$) | 26 (12.6) | 106 (41.4) | 0.0040 |
| | Mild (45-54%) | 61 (29.6) | 78 (30.5) | |
| | Moderate (31-44%) | 76 (36.9) | 42 (16.4) | |
| | Severe ($\leq 30\%$) | 43 (20.9) | 30 (11.7) | |
| Culprit Lesion | RCA | 84 (40.4) | 72 (28.1) | <0.0010 |
| | LAD | 114 (54.8) | 104 (40.6) | |
| | LCx | 10 (4.7) | 72 (28.1) | |
| | LM | 0 (0.0) | 8 (3.2) | |

On echocardiographs, NSTEMI patients had limited regional wall-motion abnormality ($p=0.0045$) compared to STEMI. Coronary angiography revealed that the left anterior descending (LAD) artery was the most commonly involved in both STEMI (54.8%) and NSTEMI patients (40.6%); however, the left circumflex (LCx) artery played a more important role in NSTEMI than in STEMI patients (28.1% vs 4.7%; $p<0.001$). With regard to short-term outcome heart failure, cardiogenic shock, arrhythmias, re-infarction and death were more in STEMI patients and it was statistically significance (Table-II).

Table-II: Short term outcome after MI

| Outcome | STEMI, n (%) | NSTEMI, n (%) | P value |
|------------------------------|---------------------|----------------------|----------------|
| Heart failure | 75 (36%) | 24 (9.3%) | 0.001 |
| Atrial fibrillation | 15 (7.2%) | 2 (0.78%) | |
| Cardiogenic shock | 35 (16.8%) | 40 (15.6%) | |
| Supraventricular Tachycardia | 5 (2.4%) | 0 (0%) | |
| Ventricular Tachycardia | 6 (2.8%) | 0 (0%) | |
| Post infarct angina | 12 (5.7%) | 38 (14.84%) | |
| Re infarction | 7 (3.3%) | 2 (0.78%) | |
| Death | 5 (2.4%) | 1 (0.40%) | |

Discussion

In this study, the mean age of STEMI and NSTEMI groups were 54.36 ± 10.18 and 51.29 ± 11 years respectively with an age range from 40 to 78 years. Majority of STEMI patients were found in 51 to 60 years of age but NSTEMI patients were mostly above 70 years. Burazeri et al⁶ found that mean age of the study subjects with STEMI was 59.1 ± 8.7 years in their study. In STEMI group male and female ratio was (4:1). In non STEMI group male and female ratio was (3:1.9).

In NSTEMI-MI patients complaints of retro sternal chest pain associated with other symptoms such as dyspnoea, nausea, abdominal pain, and syncope. However, atypical presentations are not uncommon⁷. These include stabbing chest pain, increasing dyspnoea, epigastric pain, indigestion. Atypical complaints are often observed in older patients over 75 years, in women, and in patients with diabetes, chronic kidney diseases, or dementia^{8,9}. In this study, most of NSTEMI patients presented with atypical central chest pain, abdominal pain, nausea, vomiting, sweating.

Patients who were treated with early revascularization having low (2.5%) risk for development of life-threatening arrhythmias with 80% occurring during the first 12 hours after onset of symptoms¹⁰. In this study, ST elevation was seen in all STEMI subjects and ST depression was observed in 53.2% subjects with NSTEMI. Arrhythmia including atrial fibrillation, supraventricular tachycardia, ventricular tachycardia were 12.4.% and 2% in STEMI and NSTEMI respectively. Most common echocardiographic findings of the subjects were regional wall motion abnormalities. Majority of the STEMI patients had LV dysfunction. Only 12.6% in STEMI and 41.4% in non STEMI had

>55% LV function. Coronary angiography revealed that the left anterior descending (LAD) artery was the most commonly involved artery in both STEMI (54.8%) and NSTEMI patients (40.6%); but, the left circumflex (LCx) artery played a more important role in NSTEMI (28.1%) than in STEMI patients (4.7%; $p < 0.001$). This result is consistent with Chun et al study¹¹. Recurrent post infarct angina occur in NSTEMI about 14.84% patients in this study. Effective early revascularization is beneficial for all STEMI patients. Randomized clinical trials have compared conservative treatment with invasive treatments in NSTEMI-ACS which shows that an invasive procedure reduced the incidence of myocardial infarction but not death^{12,13,14}.

In this study, 5 patients died in STEMI and 1 in NSTEMI. STEMI patients had higher hospital mortality rate than NSTEMI-ACS (7% vs 3–5%, respectively), but at 6 months mortality rates were nearly similar in both conditions (12% and 13%, respectively)¹⁵⁻¹⁷. Long-term follow-up showed that death rates were higher among NSTEMI-ACS patients than with STEMI-ACS, with a two-fold difference at 4 years¹⁸. This difference in mid-term and long-term evolution may be due to different patient profiles as NSTEMI-ACS patients were mostly older, having multiple co-morbidities, especially diabetes and renal failure. Similar study was also carried out in Bangladesh, the result of which was also consistent with this one¹⁹.

Limitations

This was a retrospective study of patients with a discharge diagnosis of STEMI or NSTEMI. In clinical practice, most of STEMI patients were managed by primary PCI who presented earlier, while NSTEMI-ACS patients received less coronary intervention. Patients enrolled in the study came from a single hospital. A regional, or even national, study should be undertaken before the results can be extrapolated to the general population.

Conclusion

At the end, it is evident that short term clinical outcome of STEMI is worse than NSTEMI. STEMI patients had higher mortality rate from acute complications within 6 months, whereas the mortality rate of NSTEMI-ACS patients were higher after 6 months. So, in clinical practice this suggests that, STEMI patients should be early revascularized to reduce mortality or morbidity. On the other hand, as most of NSTEMI-ACS patients are older and have multiple co-morbidities, so they should be counseled to increase medication adherence for secondary prevention of ischemic heart disease.

References

1. Anderson JL, Adams CD, Antman EM et al. ACC/AHA 2007 guidelines for the management of patients with unstable angina/non-ST-elevation myocardial infarction: A report of the American College of Cardiology. *Circulation* 2007; 116(7):e148-304.
2. Antman EM, Hand M, Armstrong PW et al. 2007 focused update of the ACC/AHA 2004 guidelines for the management of patients with ST-elevation myocardial infarction: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2008; 117(2):296-329.
3. Kushner FG, Hand M, Smith SC Jr et al. 2009 focused updates: ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction (updating the 2004 guideline and 2007 focused update) and ACC/AHA/SCAI guidelines on percutaneous coronary intervention (updating the 2005 guideline and 2007 focused update). *J Am Coll Cardiol* 2009; 54(23):2205-41.
4. Wagner GS, Macfarlane P, Wellens H et al. AHA/ACCF/HRS recommendations for the standardization and interpretation of the electrocardiogram. *J Am Coll Cardiol* 2009; 53(11):1003-11.
5. Van de Werf F, Bax J, Betriu A et al. Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation: The Task Force on the Management of ST-Segment Elevation Acute Myocardial Infarction of the European Society of Cardiology. *Eur Heart J* 2008; 29(23):2909-45.
6. Burazeri G, Goda A, Sulo G et al. Conventional Risk Factors and Acute Coronary Syndrome during a Period of Socioeconomic Transition: Population-based Case-control Study in Tirana, Albania. *Croat Med J* 2007; 48(2):225-33.
7. Canto JG, Fincher C, Kiefe CI et al. Atypical presentations among Medicare beneficiaries with unstable angina pectoris. *Am J Cardiol* 2002; 90:248-53.
8. Culic V, Eterovic D, Miric D et al. Symptom presentation of acute myocardial infarction: influence of sex, age, and risk factors. *Am Heart J* 2002; 144:1012-7.
9. Brieger D, Eagle KA, Goodman SG et al. Acute coronary syndrome without chest pain, an underdiagnosed and under treated high risk group: Insights from Global registry of acute coronary events. *Chest* 2004; 126(2):461-9.
10. Rahimi K, Watzlawek S, Thiele H et al. Incidence, time course, and predictors of early malignant ventricular arrhythmias after non-ST-segment elevation myocardial infarction in patients with early invasive treatment. *Eur Heart J* 2006; 27:1706-11.
11. Liu CH, Huang YC. Comparison of STEMI and NSTEMI patients in the emergency department. *JACME* 2011; 1:1-4.

12. Spacek R, Widimský P, Straka Z et al. Value of first day angiography/angioplasty in evolving Non-ST segment elevation myocardial infarction: An open multicenter randomized trial. The VINO Study. *Eur Heart J* 2002; 23:230-8.
13. Fox KA, Poole-Wilson PA, Henderson RA et al. Interventional versus conservative treatment for patients with unstable angina or non-ST-elevation myocardial infarction: The British Heart Foundation RITA 3 randomised trial. *Randomized Intervention Trial of unstable Angina. Lancet* 2002; 360(9335):743-51.
14. Hirsch A, Windhausen F, Tijssen JGP et al. Long-term outcome after an early invasive versus selective invasive treatment strategy in patients with non-ST-elevation acute coronary syndrome and elevated cardiac troponin T (the ICTUS trial): A follow-up study. *Lancet* 2007; 369:827-35.
15. Fox KA, Eagle KA, Gore JM et al. The Global Registry of Acute Coronary Events, 1999 to 2009--GRACE. *Heart* 2010; 96(14):1095-1101.
16. Savonitto S, Ardissino D, Granger CB et al. Prognostic value of the admission electrocardiogram in acute coronary syndromes. *JAMA* 1999; 281:707-13
17. Mandelzweig L, Battler A, Boyko V et al. The second Euro Heart Survey on acute coronary syndromes: Characteristics, treatment and outcome of patients with ACS in Europe and the Mediterranean Basin in 2004. *Eur Heart J* 2006; 27:2285-93.
18. Terkelsen CJ, Lassen JF, Norgaard BL et al. Mortality rates in patients with ST-elevation vs. non-ST-elevation acute myocardial infarction: Observations from an unselected cohort. *Eur Heart J* 2005; 26:18-26.
19. Rahman A, Khan MMZA, Alam MZ et al. Comparison of Short-Term Clinical Outcome in In-Hospital Patients of ST Elevation versus Non ST Elevation Myocardial Infarction. *Bangladesh Heart Journal* 2016; 31(2):70-4.